

# UNCLASSIFIED

AD NUMBER
AD852796
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; MAY 1969. Other requests shall be referred to Department of the Army, Fort Detrick, Attn: Technical Release Branch/TID, Frederick, MD 21701.
AUTHORITY
Biological Defense Research Lab ltr dtd 29 Sep 1971

THIS PAGE IS UNCLASSIFIED

AD

AD852796

TECHNICAL MANUSCRIPT 531

THE GRANULAR PNEUMONOCYTE:  
ABSENCE OF PHAGOCYTIC ACTIVITY

Charles S. Faulkner II  
John R. Esterly

MAY 1969

MAY 29 1969

DEPARTMENT OF THE ARMY  
Fort Detrick  
Frederick, Maryland

STATEMENT #2 UNCLASSIFIED  
This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of Dept. of Army, Fort Detrick, ATTN: Technical Release Branch/T10, Frederick, Maryland 21701

Reproduction of this publication in whole or in part is prohibited except with permission of the Commanding Officer, Fort Detrick, ATTN: Technical Releases Branch, Technical Information Division, Fort Detrick, Frederick, Maryland, 21701. However, DDC is authorized to reproduce the publication for United States Government purposes.

#### DDC AVAILABILITY NOTICES

Qualified requesters may obtain copies of this publication from DDC.

Foreign announcement and dissemination of this publication by DDC is not authorized.

Release or announcement to the public is not authorized.

#### DISPOSITION INSTRUCTIONS

Destroy this publication when it is no longer needed. Do not return it to the originator.

The findings in this publication are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

ACCESSION FOR	
CFSTI	WHITE SECTION <input type="checkbox"/>
DDC	BUFF SECTION <input checked="" type="checkbox"/>
UN ANNOUNCED	
JUSTIFICATION	
DISTRIBUTION/AVAILABILITY CODES	
DIST.	AVAIL. AND/OR SPECIAL
2	

DEPARTMENT OF THE ARMY  
Fort Detrick  
Frederick, Maryland 21701

TECHNICAL MANUSCRIPT 531

THE GRANULAR PNEUMONOCYTE:  
ABSENCE OF PHAGOCYTIC ACTIVITY

Charles S. Faulkner II

John R. Esterly

Pathology Division  
MEDICAL SCIENCES LABORATORIES

Project 1T061101A91A

May 1969

In conducting the research described in this report, the investigators adhered to the "Guide for Laboratory Animal Facilities and Care," as promulgated by the Committee on the Guide for Laboratory Animal Facilities and Care of the Institute of Laboratory Animal Resources, National Academy of Sciences-National Research Council.

#### ABSTRACT

Pulmonary phagocytosis has interested many investigators, but relatively few have used ultrastructural techniques for precise identification of pulmonary cells. The objective of this experiment was to determine the characteristics and variation in appearance of phagocytic cells under normal conditions and to observe the relationship between ingested particles and other cytoplasmic inclusions. A solution of India ink and polystyrene spheres was injected into the trachea of adult New Zealand rabbits. The animals were sacrificed after 45 to 60 minutes and specimens were obtained for light and electron microscopy. The particles were easily recognized in the airspaces, both free and in the cytoplasm of alveolar macrophages and occasional polymorphonuclear cells. Particles were often adjacent to granular or membranous pneumonocytes, but in no instance was ink or plastic identified within these epithelial cells, nor were particles found in interstitial or intravascular spaces. Lamellar inclusions were present in nearly all of the granular pneumonocytes (type II cell, great alveolar cell). In several samples, inclusions communicated openly with the alveolar space, but the observations do not support the hypothesis of a phagocytic origin of these cytoplasmic bodies. Phagocytosis in specimens from neonatal rabbits was not significantly different from that in the adult. These observations seem to indicate that the granular pneumonocyte is not a phagocytic cell, and that if it ever becomes phagocytic, it undergoes considerable change in its morphologic characteristics. No intermediate forms suggesting a transition from granular to phagocytic pneumonocyte have ever been observed by the authors.

# THE GRANULAR PNEUMONOCYTE: ABSENCE OF PHAGOCYTIC ACTIVITY\*

A definite alveolar epithelium was recognized and its cell types were described as a result of the magnification and resolution provided by the electron microscope.<sup>1-3</sup> There are two types of cells whose morphologic features are well known, although their origin and function are less completely understood. The membranous pneumonocyte (type I alveolar cell) is an attenuated squamous cell that lines the major portion of the alveolar surface. The granular pneumonocyte (type II alveolar cell, great alveolar cell) is more nearly cuboidal and often occupies corner positions in the alveolus. It has a vacuolated cytoplasm and electronmicrographs show characteristic lamellar inclusions that are thought by many investigators to be the precursor or storage form of pulmonary surfactant, the anti-atelectasis factor.<sup>4,5</sup> Available data suggest that both cell types are renewed by mitosis.<sup>6</sup> Nothing is known of the origin or transformation of one cell type from another, nor has a definite metabolic function been attributed to the membranous pneumonocyte.

In addition, the pulmonary septa contain capillary endothelial cells and reticuloendothelial cells. The latter appear as mononuclear leukocytes and histiocytes in septa and as alveolar macrophages in the airspaces. These cells are actively phagocytic and the majority are of bone marrow origin.<sup>7,8</sup> Their metabolism has been extensively studied, especially with respect to hydrolytic enzymes associated with phagocytosis and lysosomal activity.<sup>9,10</sup>

A recent report has suggested that lamellar inclusions in the granular pneumonocyte represent ingested rather than secretory material.<sup>11</sup> This study was undertaken to determine precisely which cells in the lung participate in phagocytosis, and to examine the cytologic features of these cells. It is a preliminary portion of a detailed study of phagocytosis and hydrolytic enzyme activity in the reticuloendothelial system.

Adult New Zealand rabbits were briefly anesthetized and 5 to 10 ml of a particulate solution were injected into the trachea. The solution was prepared by mixing, in proportions of 1:40, India ink and a 5% by volume suspension of 1-micron polystyrene spheres.\*\* The animals were sacrificed 45 to 60 min after injection. The portions of the lung into which the particulate solution had entered were easily distinguished by the India ink marker. Small fragments were fixed in phosphate-buffered 2.5% glutaraldehyde at 4 C. The specimens were subsequently osmicated, dehydrated, and embedded in Epon according to the technique of Luft.<sup>12</sup>

---

\* This report should not be used as a literature citation in material to be published in the open literature. Readers interested in referencing the information contained herein should contact the senior author to ascertain when and where it may appear in citable form.

\*\* Dow Chemicals, Inc.

Thick sections for light microscopy were stained with toluidine blue, and thin sections for electron microscopy were stained with uranyl acetate and lead citrate. Similarly processed tissues from normal-appearing areas of lung and from uninjected animals were included for comparison.


The various epithelial, interstitial, and free alveolar cell types were recognized and differentiated by location and ultrastructural features. Granular pneumonocytes were found resting directly on the alveolar basement membrane. As well as a moderate number of mitochondria, their cytoplasm contained numerous, similarly sized, rounded inclusions within which were varying amounts of a layered lipid material. The free border was formed by relatively uniform microvilli. The adjacent membranous pneumonocytes partially covered the sides of the granular pneumonocyte, and desmosomal areas of electron density were present near the alveolar side of the intercellular junctions. Alveolar macrophages, on the other hand, were not attached to the alveolar wall. Their irregular cytoplasmic projections varied considerably in size and shape and were clearly distinguished from the microvilli of the granular pneumonocyte. Multiple structures, including lysosomes, microbodies, and other larger, irregular structures containing various sorts of electron-dense material were present in the cytoplasm of the macrophage. Occasionally, the latter appeared as osmiophilic, lamellated material resembling the inclusions of the granular pneumonocyte. We have not seen any instances, however, of a cell with the other characteristics of a macrophage, containing more than one or two such bodies and causing confusion with the granular pneumonocyte. Nor have we seen any of the irregular phagosomes, which are common in the macrophage, in any of the granular pneumonocytes. In practice, the distinction between the two cells was readily apparent (Fig. 1), and no transitional forms were recognized. In only rare cases, with tangential sectioning and insufficient cytoplasmic area, was identification difficult. Plasma cells, lymphocytes, and monocytes were identified in vascular channels, and neutrophils were present both in the blood vessels and, to a lesser extent, in the airspaces.

The injected particulate material was found free in the airways and within cells (Fig. 2). Multiple carbon particles and several plastic spheres were found in 90 to 95% of the macrophages, and in all of the less frequent intra-alveolar neutrophils. Carbon and plastic were also found in very close proximity to granular pneumonocytes, but particulate matter was never seen in these or in any of the other pulmonary cell types (Fig. 3). Nor was there other evidence of phagocytosis by granular pneumonocytes. In several fields, an inclusion of a granular pneumonocyte was observed in communication with an alveolar space, but even then, with the plastic and carbon in close proximity, particles were never found within the inclusion (Fig. 4).



FIGURE 1. Adjacent Granular Pneumocyte and Alveolar Macrophage, Illustrating the Cytologic Features that Allow Differentiation between the Two Cell Types in Most Instances (see text). Enlarged from 9100X





**FIGURE 2. Alveolar Macrophage, Containing Abundant Plastic Spheres and Carbon Particles. One plastic sphere is being engulfed. 7900X**




FIGURE 1. Granular pneumonocyte, containing relatively few inclusions, and a macrophage containing plastic spheres and carbon. Carbon appears to be in contact with the surface of the granular pneumonocyte (arrow), but there is no evidence of phagocytosis by that cell. Enlarged slightly from source.




FIGURE 4. Granular Pneumonocyte, Illustrating Open Communication between the Interior of a Lamellar Inclusion and the Alveolar Space. Although carbon particles and a plastic sphere are near the opening, there is no evidence that the granular pneumonocyte is incorporating them. 13,200X

A third type of pulmonary epithelial cell (alveolar brush cell) described by Meyrick and Reid<sup>13</sup> was not recognized. Occasional granular pneumonocytes without lamellar inclusions were consistent with the description of brush cells, but they were never seen adjacent to other granular pneumonocytes, nor did they constitute even a portion of 10% of epithelial cells.<sup>13</sup>

Although it has been suggested that pulmonary epithelial cells have phagocytic capacity,<sup>14-16</sup> the only recent study with ultrastructural identification is the report of Niden. He described carbon particles in lamellar inclusions and concluded that the inclusions were a form of lysosome because of their acid phosphatase reactive membrane.<sup>11</sup> The findings in this study again demonstrate the active phagocytosis of alveolar macrophages but, within the experimental conditions, fail to confirm that the granular pneumonocyte is a phagocytic cell. Similar observations have also been reported in dogs exposed to aerosols of beryllium dust.<sup>17</sup>

Hydrolytic enzymes have been found in granular pneumonocytes by several investigations, but the observation is not necessarily evidence of lysosomal origin, since these reactions have been localized to secretory vacuoles and other cytoplasmic structures.<sup>18-21</sup> Furthermore, it has been argued that the lamellar inclusion is not phagocytosed material because it lacks a carbohydrate coat.<sup>22</sup>

The correlation of pulmonary stability and the presence of lamellar inclusions is based on a variety of circumstantial data, including appearance during gestation, experimental toxicity, and the localization of metabolic precursors to granular pneumonocytes.<sup>23-26</sup> Although precise characterization of pulmonary surfactant and its site of synthesis is lacking, these data implicate the importance of lamellar inclusions in the process, and are difficult to reconcile with their origin by phagocytosis. This study offers no additional evidence for the origin of surfactant, but it does indicate that particulate matter is not phagocytosed by the granular pneumonocytes, and it suggests that phagocytosis is an unlikely origin for lamellar inclusions.

# LITERATURE CITED

1. Low, F.N. 1952. Electron microscopy of rat lung. *Anat. Rec.* 113:437-449.
2. Karrer, H.E. 1956. The ultrastructure of mouse lung. *J. Biophys. Biochem. Cytol.* 2:241-252.
3. Policard, A.; Callet, A.; Pregermain, S. 1958. Constitution inframicroscopique et conditions de formation de certaines structures lamellaires concentriques dans des cellules pulmonaires en voie de dégénérescence expérimentale. *Compt. Rend.* 274:981-983.
4. Avery, M.E.; Said, S. 1965. Surface phenomena in lungs in health and disease. *Medicine* 44:503-526.
5. Pattle, R.E. 1965. Surface lining of lung alveoli. *Physiol. Rev.* 45:48-79.
6. Bertalanffy, F.D. 1967. Dynamics of cellular populations in the lung, p. 19-30. *In* A.A. Liebow and D.E. Smith (ed.) *The lung*. The Williams & Wilkins Company, Baltimore, Maryland.
7. Pinkett, M.O.; Cowdrey, C.R.; Nowell, P.C. 1966. Mixed hematopoietic and pulmonary origin of alveolar macrophages as demonstrated by chromosome markers. *Amer. J. Pathol.* 48:859-868.
8. Virolainen, M. 1968. Hematopoietic origin of macrophages as studied by chromosome markers in mice. *J. Exp. Med.* 127:943-952.
9. Dannenberg, A.M.; Burstone, M.S.; Walter, P.C.; Kinsley, J.W. 1963. A histochemical study of phagocytic and enzymatic functions of rabbit mononuclear and polymorphonuclear exudate cells and alveolar macrophages. *J. Cell Biol.* 17:465-486.
10. Yarborough, D.J.; Meyer, O.T.; Dannenberg, A.M.; Pearson, B. 1967. Histochemistry of macrophage hydrolases: III. Studies on  $\beta$ -galactosidase,  $\beta$ -glucuronidase and aminopeptidase with indolyl and naphthyl substrates. *J. Reticuloendothelial Soc.* 4:390-408.
11. Niden, A.H. 1967. Bronchiolar and large alveolar cell in pulmonary phospholipid metabolism. *Science* 158:1323-1324.
12. Luft, J.H. 1961. Improvements in epoxy resin embedding methods. *J. Biophys. Biochem. Cytol.* 9:409-414.
13. Meyrick, B.; Reid, L. 1968. The alveolar brush cell in rat lung: A third pneumonocyte. *J. Ultrastruct. Res.* 23:71-80.
14. Low, F.N.; Sampaio, M.M. 1957. The pulmonary alveolar epithelium as an entodermal derivative. *Anat. Rec.* 127:51-64.

15. Karrer, H.E. 1958. The ultrastructure of mouse lung: The alveolar macrophage. *J. Biophys. Biochem. Cytol.* 4:693-700.
16. Boren, H.G. 1968. Pulmonary response to inhaled carbon: A model of lung injury. *Yale J. Biol. Med.* 40:364-393.
17. Robinson, F.R.; Schaffner, F.; Trachtenberg, E. 1968. Ultrastructure of the lungs of dogs exposed to beryllium-containing dusts. *Arch. Environ. Health* 17:193-203.
18. Balis, J.U.; Conen, P.E. 1964. The role of alveolar inclusion bodies in the developing lung. *Lab. Invest.* 13:1215-1229.
19. Sorokin, S.P. 1966. A morphologic and cytochemical study on the great alveolar cell. *J. Histochem. Cytochem.* 14:884-897.
20. Goldfischer, S.; Kikkawa, Y.; Hoffman, L. 1968. The demonstration of acid hydrolase activities in the inclusion bodies of type II alveolar cells and other lysosomes in the rabbit lung. *J. Histochem. Cytochem.* 16:102-109.
21. Vatter, A.E.; Reiss, O.K.; Newman, J.K.; Lindquist, K.; Groeneboer, M. 1968. Enzymes of the lung: I. Detection of esterase with a new cytochemical method. *J. Cell Biol.* 38:80-98.
22. Kuhn, C. 1968. Cytochemistry of pulmonary alveolar epithelial cells. *Amer. J. Pathol.* 53:809-833.
23. Kikkawa, Y.; Motoyama, E.K.; Cook, C.D. 1965. The ultrastructure of the lungs of lambs. *Amer. J. Pathol.* 47:877-903.
24. Schaefer, K.E.; Avery, M.E.; Bensch, K. 1964. Time course of changes in surface tension and morphology of alveolar epithelial cells in CO<sub>2</sub>-induced hyaline membrane disease. *J. Clin. Invest.* 43:2080-2093.
25. Buckingham, S.; Heinemann, H.O.; Sommers, S.C.; McNary, W.F. 1966. Phospholipid synthesis in the large pulmonary alveolar cell. *Amer. J. Pathol.* 48:1027-1042.
26. Faulkner, C.S. 1969. The role of the granular pneumonocyte in surfactant metabolism: An autoradiographic study. *Arch. Pathol.* (in press).

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R & D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION
Department of the Army Fort Detrick, Frederick, Maryland, 21701		Unclassified
		2b. GROUP
3. REPORT TITLE		
THE GRANULAR PNEUMONOCYTE: ABSENCE OF PHAGOCYTIC ACTIVITY		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5. AUTHOR(S) (First name, middle initial, last name)		
Charles S. Faulkner II John R. Esterly		
6. REPORT DATE	7a. TOTAL NO. OF PAGES	7b. NO. OF REFS
May 1969	15	26
8a. CONTRACT OR GRANT NO.		8b. ORIGINATOR'S REPORT NUMBER(S)
a. PROJECT NO. 1T061101A91A		Technical Manuscript 531
c.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)
d.		
10. DISTRIBUTION STATEMENT		
Qualified requesters may obtain copies of this publication from DDC. Foreign announcement and dissemination of this publication by DDC is not authorized. Release or announcement to the public is not authorized.		
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY
		Department of the Army Fort Detrick, Frederick, Maryland, 21701
13. ABSTRACT		
<p>Pulmonary phagocytosis has interested many investigators, but relatively few have used ultrastructural techniques for precise identification of pulmonary cells. The objective of this experiment was to determine the characteristics and variation in appearance of phagocytic cells under normal conditions and to observe the relationship between ingested particles and other cytoplasmic inclusions. A solution of India ink and polystyrene spheres was injected into the trachea of adult New Zealand rabbits. The animals were sacrificed after 45 to 60 minutes and specimens were obtained for light and electron microscopy. The particles were easily recognized in the airspaces, both free and in the cytoplasm of alveolar macrophages and occasional polymorphonuclear cells. Particles were often adjacent to granular or membranous pneumonocytes, but in no instance was ink or plastic identified within these epithelial cells, nor were particles found in interstitial or intravascular spaces. Lamellar inclusions were present in nearly all of the granular pneumonocytes (type II cell, great alveolar cell). In several samples, inclusions communicated openly with the alveolar space, but the observations do not support the hypothesis of a phagocytic origin of these cytoplasmic bodies. Phagocytosis in specimens from neonatal rabbits was not significantly different from that in the adult. These observations seem to indicate that the granular pneumonocyte is not a phagocytic cell, and that if it ever becomes phagocytic, it undergoes considerable change in its morphologic characteristics. No intermediate forms suggesting a transition from granular to phagocytic pneumonocyte have ever been observed by the authors.</p>		
14. Key Words		
*Lung	*Alveoli pulmonis	Rabbits
*Phagocytes	Leukocytes	Epithelium

DD FORM 1473

REPLACES DD FORM 1473, 1 JAN 64, WHICH IS OBSOLETE FOR ARMY USE.

Unclassified

Security Classification